

ON THE GROWTH OF THE BERGMAN KERNEL NEAR AN INFINITE-TYPE POINT

GAUTAM BHARALI

ABSTRACT. We study diagonal estimates for the Bergman kernels of certain model domains in \mathbb{C}^2 near boundary points that are of infinite type. To do so, we need a mild structural condition on the defining functions of interest that facilitates *optimal* upper and lower bounds. This is a mild condition; unlike earlier studies of this sort, we are able to make estimates for *non-convex* pseudoconvex domains as well. This condition quantifies, in some sense, how flat a domain is at an infinite-type boundary point. In this scheme of quantification, the model domains considered below range — roughly speaking — from being “mildly infinite-type” to very flat at the infinite-type points.

1. STATEMENT OF RESULTS

Let $\Omega \subset \mathbb{C}^2$ be a pseudoconvex domain (not necessarily bounded) having a smooth boundary. Let $p \in \partial\Omega$ be a point of infinite type: by this we mean that for *each* $N \in \mathbb{Z}_+$, there exists a germ of a 1-dimensional complex-analytic variety through p whose order of contact with $\partial\Omega$ at p is at least N . If $\partial\Omega$ is not Levi-flat around p , there exist holomorphic coordinates $(z, w; V_p)$ centered at p such that

$$(1.1) \quad \Omega \cap V_p = \{(z, w) \in V_p : \Im w > F(z) + R(z, \Re w)\},$$

where F is a smooth, subharmonic, non-harmonic function defined in a neighbourhood of $z = 0$, that vanishes to infinite order at $z = 0$; $R(\cdot, 0)$ vanishes to infinite order at $z = 0$; and R is $O(|z||\Re w|, |\Re w|^2)$. Given the infinite order of vanishing of F at $z = 0$, how does one find estimates for the Bergman kernel of Ω near p ? In many cases — for instance: when $\partial\Omega \cap V_p$ is pseudoconvex of strict type, in the sense of [5], away from $p \in \partial\Omega$ — the function F in (1.1) can be extended to a global subharmonic function. In such situations, the model domain

$$(1.2) \quad \Omega_F := \{(z, w) \in \mathbb{C}^2 : \Im w > F(z)\},$$

approximates $\partial\Omega$ to infinite order along the complex-tangential directions at p . One is thus motivated to investigate estimates for the Bergman kernel for domains of the form (1.2). In this paper, we shall find estimates for the Bergman kernel of Ω_F on the diagonal as one approaches $(0, 0) \in \partial\Omega_F$. More specifically:

- We shall derive estimates that hold *not just* in a non-tangential interior cone with vertex at $(0, 0)$, but for a family of much larger approach regions that comprises regions with *arbitrarily high* orders of contact at $(0, 0)$; and

1991 *Mathematics Subject Classification*. Primary: 32A25, 32A36.

Key words and phrases. Bergman kernel, diagonal estimates, infinite type, optimal estimates. This work is supported in part by a grant from the UGC under DSA-SAP, Phase IV..

- We shall find optimal estimates for the growth of the kernel (evaluated on the diagonal of $\Omega_F \times \Omega_F$) as $(z, w) \rightarrow (0, 0)$ through any of the aforementioned approach regions.

Pointwise estimates, and a lot more, have been obtained for finite-type domains in \mathbb{C}^2 ; see for instance [2] by Diederich *et al*; [8] and [9] by Nagel *et al*; and [7] by McNeal. In [4], Kim and Lee provide some estimates on the diagonal for the Bergman kernel, as one approaches an infinite-type boundary point, for a class of convex, infinite-type domains in \mathbb{C}^2 . However, to the best of our knowledge, not even pointwise estimates are known for any reasonably general class of pseudoconvex (not necessarily convex) domains of infinite type. Determining such estimates even for model domains of the form (1.2) is not so easy. For instance, techniques analogous to the scaling methods used in the papers [7] and [9], in [1] by Boas *et al*, and in [6] by Krantz-Yu do not seem to yield optimal estimates. Another problem is that we do not know *a priori* whether Ω_F — recall that our models *do not* arise as limits of scalings of bounded domains — even has a non-trivial Bergman space. Things become tractable if we impose a simplifying condition on F :

$$(*) \left\{ \begin{array}{l} F \text{ is a radial function, i.e. } F(z) = F(|z|) \ \forall z \in \mathbb{C}, \text{ and} \\ \exists \eta > 0 \text{ such that } F(z) \geq C|z|^\eta \text{ when } |z| \geq R \text{ for some } R > 0 \text{ and } C > 0. \end{array} \right.$$

Under this condition, Ω_F has a non-trivial Bergman space; see for instance [3] by Haslinger. However, given the condition (*), we can say more: under this condition, Ω_F has a bounded realization and thus admits a localization principle for the Bergman kernel. To state this precisely, we recall that the Bergman projection for Ω is the orthogonal projection $B_\Omega : \mathbb{L}^2(\Omega) \rightarrow \mathcal{O}(\Omega) \cap \mathbb{L}^2(\Omega)$, and the Bergman kernel is the kernel representing this projection. Let us denote the Bergman kernel of Ω_F as $B_F(Z, Z')$, $(Z, Z') \in \Omega_F \times \Omega_F$. We will denote the kernel restricted to the diagonal by $K_F(z, w) := B_F((z, w), (z, w))$. We can now state

Proposition 1.1. *Let F be a C^∞ -smooth subharmonic function that vanishes to infinite order at $0 \in \mathbb{C}$ and satisfies the condition (*). Assume that the boundary of the domain $\Omega_F := \{(z, w) \in \mathbb{C}^2 : \Im w > F(z)\}$ is not Levi-flat around the origin. Then:*

- 1) *There exists an injective holomorphic map Ψ defined in a neighbourhood of $\overline{\Omega}_F$ such that $\Psi(\Omega_F)$ is a bounded pseudoconvex domain.*
- 2) *For each polydisc Δ centered at the origin, there exists a constant $\delta \equiv \delta(\Delta) > 0$ such that*

$$(1.3) \quad \delta K_{\Omega_F \cap \Delta}(z, w) \leq K_F(z, w) \ \forall (z, w) \in \Omega_F \cap (\tfrac{1}{2}\Delta).$$

Yet, does the condition (*) confer any degree of control on the decay of F near $z = 0$ that is sufficient for optimal estimates; estimates on $K_F(z, w)$ from below in particular? Even if F is radial, $K_F(z, w) \gtrsim \|(z, w)\|^{-2}$ is the best that one expects (for non-tangential approach) without any *additional* information on F . To illustrate: the condition that $(0, 0) \in \partial\Omega_F$ is of finite type facilitates optimal estimates because, with this extra information:

- one can find constants $C, \delta > 0$, and a $M \in \mathbb{Z}_+$ such that

$$(**) \quad \mathbb{B}^2(0; \delta) \cap \{(z, w) : \Im w > C|z|^{2M}\} \subset \Omega_F \cap \mathbb{B}^2(0; \delta) \\ \subset \mathbb{B}^2(0; \delta) \cap \{(z, w) : \Im w > (1/C)|z|^{2M}\};$$

- one can now make precise estimates by exploiting the simplicity of the prototypical defining function $z \mapsto |z|^{2M}$.

Some condition that enables one to work — in the spirit of $(**)$ — with easier-to-handle prototypes of F is called for if one wants optimal estimates in the infinite-type case. It turns out that we do have useful information if the infinite-type F satisfies the condition (1.4) spelt out in Theorem 1.2 below. While this condition might look rather arbitrary, it is in fact a mild restriction. It is, in some sense, a signature of F being of infinite type: a domain Ω_F satisfying (1.4) is necessarily of infinite type at $(0, 0)$. The condition (1.4) encompasses a large class of domains, ranging from the “mildly infinite type” to the very flat at $(0, 0)$ (refer to the observations following Theorem 1.2).

We need one further piece of notation. Let $f : [0, \infty) \rightarrow \mathbb{R}$ be a strictly increasing function, and let $f(0) = 0$. We define the function Λ_f as

$$\Lambda_f(x) := \begin{cases} -1/\log(f(x)), & \text{if } 0 < x < f^{-1}(1), \\ 0, & \text{if } x = 0. \end{cases}$$

We can now state our main theorem.

Theorem 1.2. *Let F be a C^∞ -smooth subharmonic function that vanishes to infinite order at $0 \in \mathbb{C}$ and satisfies the condition $(*)$. Suppose the boundary of the domain $\Omega_F := \{(z, w) \in \mathbb{C}^2 : \Im w > F(z)\}$ is not Levi-flat around the origin.*

- 1) Define f by the relation $f(|z|) = F(z)$. Then, f is a strictly increasing function on $[0, \infty)$.
- 2) Assume that F satisfies the following condition:

$$(1.4) \quad \begin{aligned} &\exists \text{ constants } B, \varepsilon_0 > 0, \text{ and a function } \chi \in \mathcal{C}([0, \varepsilon_0]; \mathbb{R}) \text{ s.t.} \\ &\chi^p \text{ is convex on } (0, \varepsilon_0) \text{ for some } p > 0, \text{ and} \\ &(1/B)\chi(x) \leq \Lambda_f(x) \leq B\chi(x) \quad \forall x \in [0, \varepsilon_0]. \end{aligned}$$

Then, for each $\alpha > 0$ and $N \in \mathbb{Z}_+$, there exists a constant $H_{N, \alpha} > 0$, which depends only on α and N ; and $C_0, C_1 > 0$, which are independent of all parameters, such that:

$$(1.5) \quad C_0(\Im w)^{-2} [f^{-1}(\Im w)]^{-2} \leq K_F(z, w) \leq C_1(\Im w)^{-2} [f^{-1}(\Im w)]^{-2} \\ \forall (z, w) \in \mathcal{A}_{\alpha, N}, \quad 0 < \Im w < H_{N, \alpha},$$

where $\mathcal{A}_{\alpha, N}$ denotes the approach region

$$\mathcal{A}_{\alpha, N} := \left\{ (z, w) \in \Omega_F : \sqrt{|z|^2 + |\Re w|^2} < \alpha(\Im w)^{1/N} \right\}.$$

- 3) Under the assumptions of (2), there exists a constant $H_0 > 0$ that is independent of all parameters such that the left-hand inequality in (1.5) in fact holds for all $(z, w) \in \Omega_F \cap \{(z, w) : \Im w < H_0\}$.

The reader might like to see examples of domains that satisfy all the hypotheses of Theorem 1.2. We discuss two examples, beginning with a very familiar example.

Example 1.3. *Estimates for the pseudoconvex domain*

$$\Omega^\beta := \{(z, w) \in \mathbb{C}^2 : \Im w > F_\beta(z)\}$$

where:

- F_β is subharmonic;
- $F_\beta(z) = \exp(-1/|z|^\beta)$, $\beta > 0$, in a neighbourhood of $z = 0$; and
- $F_\beta(z)$ grows like $|z|^2$ for $|z| \gg 1$.

We just have to check whether F satisfies the condition (1.4). There exists an $\varepsilon_0 > 0$ such that $\Lambda_f(x) = x^\beta \forall x \in [0, \varepsilon_0]$. We pick

$$p = \begin{cases} \text{any number } q \text{ such that } q\beta > 1, & \text{if } 0 < \beta \leq 1, \\ 1, & \text{if } \beta > 1. \end{cases}$$

With such a choice for p , $(\Lambda_f)^p$ itself is convex on $(0, \varepsilon_0)$. Hence, Theorem 1.2 tells us that for each $\alpha > 0$ and $N \in \mathbb{Z}_+$, there exists a constant $H_{N,\alpha} > 0$; and $C_0, C_1 > 0$, which are independent of all parameters, such that:

$$C_0 t^{-2} (\log(1/t))^{2/\beta} \leq K_{\Omega^\beta}(z, s + it) \leq C_1 t^{-2} (\log(1/t))^{2/\beta} \\ \forall (z, s + it) \in \mathcal{A}_{\alpha, N} \text{ and } 0 < t < H_{N,\alpha}. \quad \square$$

Remark 1.4. We would like to emphasize here that Λ_f is allowed to vanish to infinite order at the origin, provided it satisfies condition (1.4). So, for example, Theorem 1.2 will provide optimal growth estimates for K_F for a domain Ω_F of the form (1.2) where

- $F(z) = \exp\{-e^{1/|z|}\}$ if $z : 0 \leq |z| \leq 1/4$; and
- $F(z)$ is so defined for $|z| \geq 1/4$ that F satisfies condition (*) and Ω_F is pseudoconvex with non-Levi-flat boundary.

Domains like these are what we informally termed above as “very flat at $(0, 0)$ ”. The methods used by Kim and Lee in [4] do not seem to work for domains like these precisely because Λ_f vanishes to infinite order.

A few technical preliminaries are needed before a proof of Theorem 1.2 can be given. It would be helpful to get a sense of the key ideas of our proof. A discussion of our methodology, plus two lemmas, are presented in Section 3. The proof itself is given in Section 4. In some sense, our key technical preliminary — without which sharp lower bounds would be tricky to derive — is the proof of Proposition 1.1. This proof will form our next section.

2. THE PROOF OF PROPOSITION 1.1

Let $\eta > 0$ be as given in the condition (*). We define

$$\kappa_\eta := \begin{cases} \text{the least positive integer } \kappa \text{ such that } \kappa > 1/\eta, & \text{if } 0 < \eta \leq 1, \\ 1, & \text{if } \eta > 1. \end{cases}$$

Define the objects

$$\begin{aligned}\Psi &= (\psi_1, \psi_2) : (z, w) \mapsto \left(\frac{(2i)^{\kappa_\eta} z}{(i+w)^{\kappa_\eta}}, \frac{i-w}{i+w} \right), \\ \Pi &:= \mathbb{C} \times \{w \in \mathbb{C} : \Im w > -1\}.\end{aligned}$$

Note that $\Psi \in \mathcal{O}(\Pi; \mathbb{C}^2)$ and that Ψ is injective on Π . Define f by the relation $f(|z|) = F(z)$. Then, under our hypotheses, f is strictly increasing, whence $F(z) \geq 0 \forall z \in \mathbb{C}$. The reader is directed to Lemma 3.1 for a proof of this fact. Thus $\Omega_F \not\subseteq \Pi$, whence Ψ is injective on Ω_F .

We now claim that $\Psi(\Omega_F)$ is bounded. To see this, note that any $(z, w) \in \Omega_F$ can be written as $(z, \Re w + i(F(z) + h))$, where $h > 0$. Thus

$$(2.1) \quad |\psi_2(z, w)|^2 = \frac{(F(z) + h - 1)^2 + (\Re w)^2}{(F(z) + h + 1)^2 + (\Re w)^2} \leq 1 \quad \forall (z, w) \in \Omega_F.$$

We used the fact that $F(z) \geq 0 \forall z \in \mathbb{C}$ to deduce this estimate. Now note that

$$|\psi_1(z, w)|^2 = \frac{4^{\kappa_\eta} |z|^2}{((F(z) + h + 1)^2 + (\Re w)^2)^{\kappa_\eta}}.$$

Let $R > 0$ and $C > 0$ be exactly as given in the condition (*). Then

$$|\psi_1(z, w)|^2 \leq \frac{4^{\kappa_\eta} R^2}{(F(z) + h + 1)^{2\kappa_\eta}} \leq 4^{\kappa_\eta} R^2 \quad \forall (z, w) \in \Omega_F \text{ and } |z| \leq R.$$

On the other hand

$$|\psi_1(z, w)|^2 \leq \frac{4^{\kappa_\eta} |z|^2}{(F(z))^{2\kappa_\eta}} \leq \frac{4^{\kappa_\eta} |z|^2}{C^{2\kappa_\eta} |z|^{2\eta\kappa_\eta}} \quad \forall (z, w) \in \Omega_F \text{ and } |z| \geq R.$$

From the last two inequalities, we conclude that

$$(2.2) \quad |\psi_1(z, w)|^2 \leq \max \left\{ 4^{\kappa_\eta} R^2, \left(\frac{4}{C^2} \right)^{\kappa_\eta} R^{-2(\eta\kappa_\eta - 1)} \right\} \quad \forall (z, w) \in \Omega_F.$$

From (2.1) and (2.2), our claim, and hence Part (1), follows.

To demonstrate Part (2), we will need a localization principle established by Ohsawa:

Localization Lemma (Ohsawa, [10]) *Let D be a bounded pseudoconvex domain in \mathbb{C}^n , p be a boundary point, and $V \Subset U$ be two open neighbourhoods of p . Then, there is a constant $\delta \equiv \delta(U, V) > 0$ such that*

$$\delta K_{D \cap U}(Z) \leq K_D(Z) \quad \forall Z \in D \cap V.$$

Substituting

$$\begin{aligned}D &= \Psi(\Omega_F), & p &= (0, 1), \\ D \cap U &= \Psi(\Omega_F \cap \Delta), & D \cap V &= \Psi(\Omega_F \cap (\tfrac{1}{2}\Delta))\end{aligned}$$

into the localization lemma, we conclude that there exists a $\delta \equiv \delta(\Delta) > 0$ such that (here G_F stands for $\Psi(\Omega_F)$)

$$(2.3) \quad \delta K_{G_F \cap U}(\Psi(z, w)) \leq K_{G_F}(\Psi(z, w)) \quad \forall (z, w) \in \Omega_F \cap (\tfrac{1}{2}\Delta).$$

Recall, however, the transformation rule for the Bergman kernel:

$$K_{\Omega^j}(z, w) = |\text{Jac}_{\mathbb{C}}(\Psi)(z, w)|^2 K_{\Psi(\Omega^j)}(\Psi(z, w)) \quad \forall (z, w) \in \Omega^j, \quad j = 1, 2,$$

where, in the present case, $\Omega^1 = \Omega_F$ and $\Omega^2 = \Omega_F \cap \Delta$. Applying this to (2.3) gives us the inequality (1.3). \square

3. PRELIMINARY REMARKS AND LEMMAS

The idea behind the upper bound in (1.5) is quite standard. Given a point $(z_0, w_0) \in \Omega_F$, the quantity $K_F(z_0, w_0)$ is dominated by the reciprocal of the volume of the largest polydisc centered at (z_0, w_0) that is contained in Ω_F . The volume of this polydisc will be influenced by the curvature of $\partial\Omega_F$ at the point on $\partial\Omega_F$ that is closest to (z_0, w_0) . However, if (z_0, w_0) is confined to any of the approach regions $\mathcal{A}_{\alpha, N}$, then this volume is controlled by the boundary geometry at $(0, 0) \in \partial\Omega_F$. That one has this control for any approach region $\mathcal{A}_{\alpha, N}$ — regardless of α and N — is a consequence of the fact that $(0, 0)$ is of infinite type.

The derivation of the lower bound in (1.5) relies on the construction of a suitable square-integrable holomorphic function. In this construction, we are aided by the localization principle stated in Proposition 1.1. The three main ingredients in the derivation of the lower bound are:

- i) We choose a suitable polydisc Δ centered at $(0, 0)$ and estimate $K_{\Omega_F \cap \Delta}(z, w)$ for $(z, w) \in \mathcal{A}_{\alpha, N} \cap (\frac{1}{2}\Delta)$. We rely on the fact that $K_{\Omega_F \cap \Delta}(z, w)$ is given by

$$K_{\Omega_F \cap \Delta}(z, w) = \sup \left\{ \frac{|\phi(z, w)|^2}{\|\phi\|_{\mathbb{L}^2(\Omega_F \cap \Delta)}^2} : \phi \in A^2(\Omega_F \cap \Delta) \right\}.$$

- ii) To obtain a lower bound, we select a suitable function $\phi_t \in A^2(\Omega_F \cap \Delta)$ and estimate $\|\phi_t\|_{A^2}^2$, $t > 0$. This reduces finding a lower bound for $K_{\Omega_F \cap \Delta}(z, s + it)$, $(z, s + it) \in \mathcal{A}_{\alpha, N} \cap (\frac{1}{2}\Delta)$, to estimating an integral over a region in \mathbb{R}^4 whose boundaries are determined by the function f .
- iii) The difficult issue is to find the *desired bound in terms of t* for the latter integral. The condition (1.4) is used to break up the aforementioned region of integration into sub-domains on which the integral admits the desired estimate.

We now present two lemmas that will be necessary to complete the proof of Theorem 1.2. Lemma 3.1 constitutes the proof of Part (1) of Theorem 1.2.

Lemma 3.1. *Let F be a smooth subharmonic function on \mathbb{C} such that $F(0) = 0$ and F is radial. Define f by the relation $f(|z|) = F(z)$, and write $\Omega_F := \{(z, w) \in \mathbb{C}^2 : \Im w > F(z)\}$. Assume that Ω_F is not Levi-flat in a neighbourhood of $(0, 0)$. Then f is a strictly increasing function on $[0, \infty)$.*

Proof. Suppose there exist $r_1 < r_2$, with $r_1, r_2 \in [0, \infty)$, such that $f(r_1) \geq f(r_2)$. Then, by our hypothesis on F

$$\sup_{z \in \partial D(0; r_2)} F(z) = f(r_2) \leq f(r_1).$$

By the Maximum Principle, therefore, $F|_{D(0; r_2)} \equiv 0$. But then, this would imply that the portion $\partial\Omega_F \cap \mathbb{B}^2(0; r_2)$ of $\partial\Omega_F$ is Levi-flat; i.e. a contradiction. Hence f is strictly increasing. \square

Lemma 3.2. *Let F and Ω_F have all the properties listed in Lemma 3.1. Let Λ_f satisfy the condition (1.4), and let (since, in view of Lemma 3.1, f is increasing) $G_f := \Lambda_f^{-1}$. Then, there exist constants $T > 0$ and $K > 0$ such that*

$$(3.1) \quad 0 < G_f(2t)^2 - G_f(t)^2 \leq KG_f(t)^2 \quad \forall t \in (0, T).$$

Proof. We just have to show that there exist $T > 0$ and $M > 0$ such that

$$(3.2) \quad 0 < G_f(2t) - G_f(t) \leq MG_f(t) \quad \forall t \in (0, T).$$

If we could show this, then it would follow that

$$G_f(2t)^2 - G_f(t)^2 \leq M(M+2)G_f(t)^2 \quad \forall t \in (0, T).$$

To proceed further, we need the following:

Fact. *Let g be a continuous, strictly increasing function on $[0, R]$ satisfying $g(0) = 0$, and assume g^p is convex on $(0, R)$ for some $p > 0$. Define $G := g^{-1}$, and let $B > 1$. Then:*

$$(3.3) \quad \frac{G(Bt)}{G(t)} \leq B^p \quad \forall t \in (0, g(R)/B).$$

To verify this fact, set $\Phi := (g^p)^{-1}$. Then:

$$(3.4) \quad \Phi(t) = G(t^{1/p}) \quad \forall t \in [0, g(R)^p],$$

By hypothesis, Φ is concave on $(0, g(R)^p)$. But since Φ is also continuous,

$$\frac{\Phi(B^p t^p)}{B^p t^p} \leq \frac{\Phi(t^p)}{t^p} \quad \forall t \in (0, g(R)/B).$$

The above fact now follows simply by rearranging the terms in the above inequality, and applying (3.4).

Now let ε_0 , B , χ , and p be as in (1.4). Let us also define

$$\begin{aligned} \varkappa_0 &:= (\chi)^{-1} : [0, \chi(\varepsilon_0)] \longrightarrow \mathbb{R} \\ \varkappa_1 &:= (B\chi)^{-1} : [0, B\chi(\varepsilon_0)] \longrightarrow \mathbb{R} \\ \varkappa_2 &:= [(1/B)\chi]^{-1} : [0, (1/B)\chi(\varepsilon_0)] \longrightarrow \mathbb{R} \end{aligned}$$

Since Λ_f and χ are strictly increasing (in view of Lemma 3.1), condition (1.4) implies that:

$$\varkappa_1(t) \leq G_f(t) \leq \varkappa_2(t) \quad \forall t \in [0, T_1],$$

where $T_1 := (1/B)\chi(\varepsilon_0)$. Therefore, we get

$$(3.5) \quad \frac{G_f(2t)}{G_f(t)} \leq \frac{\varkappa_2(2t)}{\varkappa_1(t)} = \frac{\varkappa_2(2t)}{\varkappa_2(t)} \frac{\varkappa_2(t)}{\varkappa_1(t)} \quad \forall t \in (0, T_1/2).$$

We now note that

$$\varkappa_1(t) = \varkappa_0(B^{-1}t), \quad \varkappa_2(t) = \varkappa_0(Bt) \quad \forall t \in [0, T_1].$$

Given this last piece of information, we can apply the inequality (3.3) to the ratios on the right-hand side of (3.5). Set $T := \min(T_1/2, (1/B)T_1)$. Then,

$$\frac{G_f(2t)}{G_f(t)} \leq (2B^2)^p \quad \forall t \in (0, T).$$

From this, the estimate (3.2) clearly follows if we take $M = (2B^2)^p - 1$. Hence, by our earlier remarks, the result follows. \square

4. THE PROOF OF THEOREM 1.2

Part (1) of Theorem 1.2 has already been established in Lemma 3.1. Therefore, f^{-1} is a well-defined function. Observe that

$$(4.1) \quad f^{-1}(t) = G_f \left(\frac{1}{\log(1/t)} \right), \quad 0 < t < 1.$$

Let $R > 0$ be so small that

$$f^{-1}(3t/4) = G_f \left(\frac{1}{\log(4/3) + \log(1/t)} \right) \geq G_f \left(\frac{1}{2 \log(1/t)} \right) \quad \forall t \in (0, R).$$

Let M and T be as given by (3.2) above. Shrinking $R > 0$ if necessary so that $0 < 1/2 \log(1/t) < T \quad \forall t \in (0, R)$, we get

$$(4.2) \quad \frac{f^{-1}(3t/4)}{f^{-1}(t)} \geq \frac{G_f(1/2 \log(t^{-1}))}{G_f(1/\log(t^{-1}))} \geq (M+1)^{-1} \quad \forall t \in [0, R].$$

Write $\mu := (M+1)^{-1}$. We are given $\alpha > 0$ and $N \in \mathbb{Z}_+$. Since $f(x)$ vanishes to infinite order at $x = 0$, there exists a $H_{N,\alpha} > 0$ such that $R \geq H_{N,\alpha}$ and

$$(4.3) \quad \alpha t^{1/N} \leq \frac{\mu}{4} f^{-1}(t) \quad \forall t \in [0, H_{N,\alpha}].$$

From (4.2) and (4.3), we see that

$$|z| + \frac{\mu}{2} f^{-1}(t) < f^{-1}(3t/4) \quad \forall z : 0 \leq |z| < \alpha t^{1/N}, \quad 0 < t < H_{N,\alpha},$$

whence the polydisc

$$(4.4) \quad \Delta(z, t) := \mathbb{D} \left(z; \frac{\mu}{2} f^{-1}(t) \right) \times \mathbb{D}(it; t/4) \subset \Omega_F$$

$$\forall z : 0 \leq |z| < \alpha t^{1/N}, \quad 0 < t < H_{N,\alpha}.$$

Now note that that translations $T_s : (z, w) \mapsto (z, s + w)$, $s \in \mathbb{R}$, are all automorphisms of Ω_F . Thus, by the transformation rule for the Bergman kernel, and by monotonicity, we get

$$\begin{aligned} K_F(z, s + it) &= K_F(z, it) \\ &\leq K_{\Delta(z,t)}(z, it) \\ &= \frac{1}{\text{vol}(\Delta(z, t))} \quad \forall (z, s + it) \in \mathcal{A}_{\alpha, N}, \quad 0 < t < H_{N,\alpha}. \end{aligned}$$

The last equality follows from the fact that $\Delta(z, t)$ is a Reinhardt domain centered at (z, t) . Hence, we have one half of the estimate (1.5):

$$(4.5) \quad K_F(z, w) \leq C_1 (\Im w)^{-2} [f^{-1}(\Im w)]^{-2} \quad \forall (z, w) \in \mathcal{A}_{\alpha, N},$$

$$0 < \Im w < H_{N,\alpha},$$

where $C_1 = 64/\mu^2 \pi^2$.

We will now derive a lower bound. We set $A := \min(f^{-1}(1), 1)$. For the remainder of this proof, Δ will denote the polydisc $\mathbb{D}(0; A) \times \mathbb{D}(0; 1)$. In view of the inequality (1.3) of Proposition 1.1, it suffices to find a lower bound for $K_{\Omega_F \cap \Delta}(z, w)$ for $(z, w) \in (\frac{1}{2}\Delta)$. It is well known that

$$(4.6) \quad K_{\Omega_F \cap \Delta}(z, w) = \sup_{\phi \in A^2(\Omega_F \cap \Delta)} \frac{|\phi(z, w)|^2}{\|\phi\|_{\mathbb{L}^2(\Omega_F \cap \Delta)}^2}.$$

Once again, we use the fact that the translations $T_u : (z, w) \mapsto (z, u + w)$, $u \in \mathbb{R}$, are all automorphisms of Ω_F , whence

$$(4.7) \quad K_F(z, s + it) = K_F(z, (u + s) + it) \quad \forall (z, s + it) \in \Omega_F \text{ and } \forall u \in \mathbb{R}.$$

Set $\phi_t(z, w) := -4t^2/(w + it)^2$, $t > 0$. Then, from the localization principle (1.3), and from (4.6) and (4.7), we get

$$(4.8) \quad \begin{aligned} K_F(z, s + it) &\geq \delta K_{\Omega_F \cap \Delta}(z, it) \\ &\geq \frac{\delta}{\|\phi_t\|_{\mathbb{L}^2(\Omega_F \cap \Delta)}^2} \quad \forall (z, t) \in (\tfrac{1}{2}\Delta). \end{aligned}$$

Let us write $w = u + iv$. We leave the reader to verify that we can apply Fubini's theorem wherever necessary in the following computation:

$$\begin{aligned} \|\phi_t\|_{\mathbb{L}^2(\Omega_F \cap \Delta)}^2 &= \int_{|z| < A} \int_{-1}^1 \int_{F(z)}^{\sqrt{1-u^2}} \frac{16t^4}{|u + i(v+t)|^4} dv du dA(z) \\ &\leq 16t^4 \int_{|z| < A} \int_{F(z)}^{\infty} \int_{-1}^1 (v+t)^{-4} \left(1 + \left(\frac{u}{v+t}\right)^2\right)^{-2} du dv dA(z) \\ &\leq 8t^4 \left(\int_{\mathbb{R}} \frac{dX}{(1+X^2)^2} \right) \int_{|z| < A} (t + F(z))^{-2} dA(z) \\ &= Ct^4 \int_0^A \frac{r}{(t + f(r))^2} dr, \end{aligned}$$

where $C > 0$ is a universal constant. In what follows, we shall denote $f^{-1}(s)$ by R_s . By equation (4.1) we have

$$(4.9) \quad R_{\sqrt{t}} = G_f \left(\frac{2}{\log(1/t)} \right), \quad 0 < t < 1.$$

We break up the interval of integration of the last integral into three sub-intervals to compute:

$$\begin{aligned}
\|\phi_t\|_{\mathbb{L}^2(\Omega_F \cap \Delta)}^2 &= Ct^4 \left(\int_0^{R_t} + \int_{R_t}^{R_{\sqrt{t}}} + \int_{R_{\sqrt{t}}}^A \frac{r}{(t+f(r))^2} dr \right) \\
&\leq Ct^4 \int_0^{R_t} \frac{r}{t^2} dr + Ct^4 \left(\int_{R_t}^{R_{\sqrt{t}}} + \int_{R_{\sqrt{t}}}^A \frac{r}{4tf(r)} dr \right) \\
&\leq \frac{C}{2} t^2 (R_t)^2 + \frac{C}{4} t^2 \int_{R_t}^{R_{\sqrt{t}}} r dr + \frac{C}{4} t^{5/2} A(A - R_{\sqrt{t}}) \\
(4.10) \quad &\leq \frac{C}{2} t^2 (R_t)^2 + \frac{C}{4} t^{5/2} A(A - R_{\sqrt{t}}) \\
&\quad + \frac{C}{8} t^2 \left(G_f^2 \left(\frac{2}{\log(1/t)} \right) - G_f^2 \left(\frac{1}{\log(1/t)} \right) \right), \quad 0 < t < 1.
\end{aligned}$$

We used the relation (4.9) in the estimate for the middle integral above.

We now apply Lemma 3.2 to the third term in (4.10). Let $T > 0$ and $K > 0$ be defined as given by Lemma 3.2. Let H_0 be so small that $1/\log(t^{-1}) < T \forall t \in (0, H_0)$, and so that the second inequality below holds true:

$$\begin{aligned}
(4.11) \quad \|\phi_t\|_{\mathbb{L}^2(\Omega_F \cap \Delta)}^2 &\leq \frac{C}{2} \left(1 + \frac{K}{4} \right) t^2 (R_t)^2 + \frac{C}{4} t^{5/2} \\
&\leq C(1 + K/4) t^2 (R_t)^2 \quad \forall t \in (0, H_0).
\end{aligned}$$

Since $f(x)$ vanishes to infinite order at $x = 0$, we can lower H_0 — and this is independent of parameters like $\alpha > 0$ and $N \in \mathbb{Z}_+$ — so that the first term of the first inequality above dominates the second for $t \in (0, H_0)$, giving us (4.11). Lowering the value of H_0 further if necessary, we also ensure:

$$\Omega_F \cap \{(z, w) : \Im w < H_0\} \subset \Omega_F \cap \Delta.$$

From (4.8) and (4.11), we conclude that there exists a constant C_0 , which is independent of all parameters, such that

$$C_0 (\Im w)^{-2} [f^{-1}(\Im w)]^{-2} \leq K_F(z, w) \quad \forall (z, w) \in \Omega_F \cap \{(z, w) : \Im w < H_0\}.$$

This establishes Part (3) of our theorem. As a special case, we get the lower bound on $K_F(z, w)$ in the estimate (1.5). Along with (4.5), this establishes Part (2) of our theorem. \square

Acknowledgements. First, and foremost, I thank Alexander Nagel for his interest in an earlier version of this article. My discussion with him led to the simplification of several arguments in that version; now the arXiv preprint [arXiv:0708.2894v1](https://arxiv.org/abs/0708.2894v1). I also thank all my colleagues who noticed this preprint and offered their comments during my stay at the Institut Mittag-Leffler during the Special Semester on complex analysis in several variables. The hospitality of the Institut Mittag-Leffler is greatly appreciated. Finally, I thank the anonymous referee of this article for his/her valuable suggestions.

REFERENCES

- [1] H.P. Boas, E.J. Straube, J. Yu, *Boundary limits of the Bergman kernel and metric*, Michigan Math. J. **42** (1995), 449-461.
- [2] K. Diederich, G. Herbort and T. Ohsawa, *The Bergman kernel on uniformly extendable pseudoconvex domains*, Math. Ann. **273** (1986), 471-478.
- [3] F. Haslinger, *Bergman and Hardy spaces on model domains*, Illinois J. Math. **42** (1998), 458-469.
- [4] K.T. Kim and S. Lee, *Asymptotic behavior of the Bergman kernel and associated invariants in certain infinite-type pseudoconvex domains*, Forum Math. **14** (2002), 775-795.
- [5] J.J. Kohn, *Boundary behavior of $\bar{\partial}$ on weakly pseudoconvex manifolds of dimension two*, J. Diff. Geom. **6** (1972), 523-542.
- [6] S.G. Krantz and J. Yu, *On the Bergman invariant and curvatures of the Bergman metric*, Illinois J. Math. **40** (1996), 226-244.
- [7] J.D. McNeal, *Boundary behavior of the Bergman kernel function in \mathbb{C}^2* , Duke Math. J. **58** (1989), 499-512.
- [8] A. Nagel, J.-P. Rosay, E.M. Stein, and S. Wainger, *Estimates for the Bergman and Szegő kernels in certain weakly pseudoconvex domains*, Bull. Amer. Math. Soc. (N.S.) **18** (1988), 55-59.
- [9] A. Nagel, J.-P. Rosay, E.M. Stein, and S. Wainger, *Estimates for the Bergman and Szegő kernels in \mathbb{C}^2* , Ann. of Math. (2) **129** (1989), 113-149.
- [10] T. Ohsawa, *Boundary behavior of the Bergman kernel function on pseudoconvex domains*, Publ. Res. Inst. Math. Sci. **20** (1984), 897-902.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE – 560 012
E-mail address: bharali@math.iisc.ernet.in